OTC Allergy reliever dosing directions for adults and children 12 years and older

Cetirizine Hydrochloride	AMOUNT (Active ingredient(s))	DOSE & FREQUENCY (Adult use only: 12 years and older)	DAILY LIMIT
Cetirizine hydrochloride regular strength for example REACTINE® Regular Strength	Cetirizine hydrochloride: 5 mg per tablet	1 or 2 tablets every 24 hours Adults 65+ years: 1 tablet every 24 hours	Do NOT take more than 2 tablets in 24 hours Adults 65+ years: 1 tablet every 24 hours
Cetirizine hydrochloride extra strength for example REACTINE® Extra Strength or REACTINE® Extra Strength Rapid Dissolve	Diphenhydramine hydrochloride: 10 mg per tablet (or rapid dissolve tablet)	1 tablet (or rapid dissolve tablet) every 24 hours Adults 65+ years: USE REACTINE® Regular Strength 5 mg	Do NOT take more than 1 tablet (or rapid dissolve tablet) in 24 hours Adults 65+ years: USE REACTINE® Regular Strength 5 mg
Cetirizine hydrochloride with Pseudoephedrine hydrochloride** for example REACTINE® Complete Sinus + Allergy	Cetirizine hydrochloride: 5 mg per tablet PLUS Pseudoephedrine hydrochloride**: 120 mg per tablet	1 tablet every 12 hours Adults 65+ years: 1 tablet every 24 hours	Do NOT take more than 2 tablets in 24 hours Adults 65+ years: Do not take more than 1 tablet every 24 hours
Cetirizine hydrochloride oral solution for example REACTINE® Liquid	Cetirizine hydrochloride: 5 mg/5 mL oral solution	5 mL or 10 mL once every 24 hours Adults 65+ years: 5 mL once every 24 hours	Do NOT take more than 10 mL in 24 hours Adults 65+ years: Do not take more than 5 mL in 24 hours

What you can do

- If provided, use only the dosing device that comes with your medicine
- Take ONLY 1 medicine that contains the same kind of medicinal ingredient at a time

Why?

If you take more than 1 medicine with the same kind of medicinal ingredient, it can increase your chance of harmful side effects

It is important to always read and follow the label. If you have any questions, ask your healthcare professional which product may be right tor you



OTC Allergy reliever dosing directions for adults and children 12 years and older

Diphenhydramine Hydrochloride	AMOUNT (Active ingredient(s))	DOSE & FREQUENCY (Adult use only: 12 years and older)	DAILY LIMIT
Diphenhydramine hydrochloride regular strength for example BENADRYL* Allergy	Diphenhydramine hydrochloride: 25 mg per caplet	1-2 caplets every 4-6 hours	Do NOT take more than 8 caplets in 24 hours Do NOT use with any other product containing diphenhydramine, even one used on skin
Diphenhydramine hydrochloride extra strength for example Extra Strength BENADRYL* Allergy	Diphenhydramine hydrochloride: 50 mg per caplet	1 caplets every 4-6 hours	Do NOT take more than 4 caplets in 24 hours Do NOT use with any other product containing diphenhydramine, even one used on skin
Diphenhydramine hydrochloride with Acetaminophen" and Pseudoephedrine hydrochloride" for example BENADRYL® Allergy, Sinus & Pain Total	Diphenhydramine hydrochloride: 25 mg per caplet PLUS Acetaminophen (extra strength)**: 500 mg per caplet PLUS Pseudoephedrine hydrochloride*: 30 mg per caplet	2 caplets every 6 hours	Do NOT take more than 8 caplets in 24 hours Do not with other products containing acetaminophen or other cough/cold or pain/ fever medications. Do NOT use with any other product containing diphenhydramine, even one used on skin
Diphenhydramine hydrochloride elixir for example BENADRYL® Elixir	Diphenhydramine hydrochloride: 1 teaspoonful (12.5 mg/5 mL)	2-4 teaspoons (10-20 mL) every 4-6 hours	Do NOT exceed 16 teaspoons (80 mL) in 24 hours Do NOT use with any other product containing diphenhydramine, even one used on skin
Diphenhydramine hydrochloride extra strength for nighttime for example Extra Strength BENADRYL® Nighttime	Diphenhydramine hydrochloride: 50 mg per caplet	1 caplet at bedtime	Take 1 caplet at bedtime if needed or as directed by a doctor Do NOT use with any other product containing diphenhydramine, even one used on skin

What you can do

- If provided, use only the dosing device that comes with your medicine
- · Take ONLY 1 medicine that contains the same kind of medicinal ingredient at a time

Whv?

If you take more than 1 medicine with the same kind of medicinal ingredient, it can increase your chance of harmful side effects

It is important to always read and follow the label. If you have any questions, ask your healthcare professional which product may be right tor you



OTC Allergy reliever dosing directions for adults and children 12 years and older

Loratadine	AMOUNT (Active ingredient(s))	DOSE & FREQUENCY (Adult use only: 12 years and older)	DAILY LIMIT
Loratadine for example CLARITIN® Allergy 24 Hour	Loratadine: 10 mg per tablet	1 tablet every 24 hours	Do NOT take more than 1 tablet in 24 hours
Loratadine orally disintegrating tablets for example CLARITIN® Rapid Dissolve 24 Hour Tablets	Loratadine: 10 mg per tablet	1 caplets every 24 hours	Do NOT take more than 1 tablet in 24 hours
Loratadine Capsules for example CLARITIN* Liquid Capsules 24 Hour	Loratadine: 10 mg per capsule	2 caplets every 24 hours	Do NOT take more than 1 capsule in 24 hours

Desloratadine	AMOUNT (Active ingredient(s))	DOSE & FREQUENCY (Adult use only: 12 years and older)	DAILY LIMIT
Desioratadine for example CLARITIN® Allergy 24 Hour	Desloratadine: 5 mg per tablet	1 tablet every 24 hours	Do NOT take more than 1 tablet in 24 hours
Desioratadine with pseudoephedrine sulfate" for example AERIUS® DoubleAction 12 Hour	Desloratadine: 2.5 mg per tablet PLUS Pseudoephedrine sulfate** 120 mg per tablet	1 caplets every 12 hours	Do NOT take more than 2 tablet in 24 hours

Fluticasone	AMOUNT (Active ingredient)	DOSE & FREQUENCY (Adult use only: 18 years and older)	DAILY LIMIT
Fluticasonepropionate (corticosteroid) nasal spray for example Flonase® Allergy Relief	Fluticasone propionate (corticosteroid): 50 mcg per spray	Week 1 Use 2 sprays in each nostril once daily, preferably in the morning. Week 2 through 3 months Use 1 spray in each nostril once daily, if your symptoms are under control. After 3 months of daily use Ask your doctor if you can keep using this product.	Week 1 No more than 2 sprays in each nostril daily. Week 2 through 3 months No more than 1 spray in each nostril daily. After 3 months of daily use Ask your doctor if you can keep using this product and what the daily limit may be.

What you can do

- If provided, use only the dosing device that comes with your medicine
- Take ONLY 1 medicine that contains the same kind of medicinal ingredient at a time

Why?

If you take more than 1 medicine with the same kind of medicinal ingredient, it can increase your chance of harmful side effects

It is important to always read and follow the label. If you have any questions, ask your healthcare professional which product may be right tor you

